Post-Operative Instructions for Ankle/Foot Surgery

□ Nimish R. Kadakia, M.D.□ Steven A. Becker, M.D.	□ Robert A. Baird, M.D.□ Floyd G. Shon, M.D.
PROCEDURE ☐ Ankle Arthroscopy ☐ Ligament Reconstruction ☐ Ankle Fracture ☐ Achilles Tendon Repair ☐ Other	 ☐ Microfracture – Cartilage Injury ☐ Tendon Reconstruction/Repair ☐ Foot Fracture ☐ Chrondroplasty – Cartilage Wear
 DRESSING ◆ Dressing can become blood stained; this is not unusual. ◆ Sutures/staples will be removed in the office. ◆ If you feel the bleeding is excessive, call the Doctor. □ You may remove dressing in 24 hours. □ Cover with gauze or ace wrap. □ OK to loosen splint if needed – please re-wrap □ Keep dressing/splint on until office visit 	 SHOWERING ☐ It is OK to shower after 24 hours. ♦ Please keep wounds dry. ♦ Wrap with plastic wrap. ♦ Redress wound after shower.
ACTIVITY ◆ Elevate operative ankle/foot above level of heart. ◆ Ice to operative ankle/foot 20 minutes/hour as tolerated. ◆ Bend and straighten knee 5-10 times per hour as tolerated. ◆ Rest, Ice, Elevation, Compression. WEIGHTBEARING ◆ You may bear % body weight on operative leg ◆ Use crutches as needed. ◆ If provided with knee brace, use for walking and sleeping. ◆ OK to remove brace while sitting.	 CALL THE DOCTOR IF YOU EXPERIENCE ◆ Persistent or abnormal bleeding. ◆ Fever or chills. ◆ Temperature greater than 101. ◆ Severe and uncontrolled pain or swelling. ◆ Medication problems. ◆ Abnormal wound drainage
 MEDICATIONS ♦ Narcotic pain medicine can make you nauseous. ♦ Take medications with food. ♦ Do NOT drive while taking pain meds. ♦ Call if pain is not well controlled. OTHER:	FOLLOW-UP APPOINTMENT
<u>+</u>	