## Post-Operative Instructions for Elbow/Hand/Wrist Surgery

| <ul><li>□ Nimish R. Kadakia, M.D.</li><li>□ Steven A. Becker, M.D.</li></ul> | <ul><li>□ Robert A. Baird, M.D.</li><li>□ Floyd G. Shon, M.D.</li></ul> |
|--|---|
| <u>PROCEDURE</u>   |   |
| □ Wrist Arthroscopy  | ☐ Carpal Tunnel Release   |
| ☐ Ligament Reconstruction  | ☐ Trigger Finger Release  |
| □ Wrist Fracture   | □ Tendon Repair   |
| ☐ Hand Fracture  | ☐ Joint Reconstruction/Arthritis  |
| ☐ Elbow Fracture   | ☐ Tennis/Golfers Elbow Repair   |
| ☐ Elbow Arthroscopy  | ☐ Biceps Tendon Repair  |
| □ Other  |   |
| <u>DRESSING</u>  | SHOWERING   |
| ♦ Dressing can become blood stained; this is not unusual.                    | ☐ It is OK to shower after 24 hours.                                    |
| ♦ If you feel the bleeding is excessive, call the Doctor.                    | ♦ Please keep wounds dry  |
| ♦ Sutures/staples will be removed in the office.                             | ♦ Wrap with plastic wrap  |
| ☐ You may remove dressing in 24 hours.                                       | ♦ Redress wound after shower.   |
| □ Cover with gauze or ace wrap.  |   |
| ☐ Use sling as directed  |   |
| ☐ OK to loosen splint if needed – please re-wrap                             |   |
| ☐ May remove provisional splint after days                                   |   |
| ☐ Do not remove splint/dressing until office visit                           |   |
| <u>ACTIVITY</u>  | CALL THE DOCTOR IF YOU  |
| ♦ Elevate operative wrist/hand above level of heart.                         | <u>EXPERIENCE</u>   |
| ♦ Ice to operative Elbow/wrist/hand 20 minutes/hour as tolerated.            | <u> </u>  |
| ☐ Bend and straighten elbow 5-10 times per hour as tolerated.                | ♦ Fever or chills.  |
| ☐ Maintain use of sling and splint until follow-up                           | ♦ Medication problems.  |
| ♦ Rest, Ice, Elevation, Compression.   | ◆ Temperature greater than 101.   |
|  | ♦ Abnormal wound drainage   |
|  | • Severe and uncontrolled pain or swelling.                             |
|  | ♦ Medication problems.  |
| <u>MEDICATIONS</u>   | FOLLOW-UP APPOINTMENT   |
| ♦ Narcotic pain medicine can make you nauseous.                              | ♦ 24-48 hours   |
| ◆ Take medications with food.  | ♦ 7-10 days   |
| ◆ Do NOT drive while taking pain meds.                                       | ◆ Call <u>949-727-3636</u> to schedule                                  |
| ♦ Call if pain is not well controlled.                                       |   |
| OTHER:   |   |
| <b>\</b>   |   |
| •  |   |